

# JUMP OF MILNOR NUMBERS

ARNAUD BODIN

**ABSTRACT.** In this note we study a problem of A'Campo about the minimal non-zero difference between the Milnor numbers of a germ of plane curve and one of its deformation.

## 1. PROBLEM OF THE JUMP (A'CAMPO)

Let  $f_0 : (\mathbb{C}^n, 0) \longrightarrow (\mathbb{C}, 0)$  be an analytic germ of isolated singularity. A *deformation* of  $f_0$  is a family  $(f_s)_{s \in [0,1]}$  of germs of isolated singularities such that the coefficients are analytic functions of  $s \in [0, 1]$ .

The *jump of the family*  $(f_s)$  is

$$\mu(f_0) - \mu(f_s), \quad 0 < s \ll 1,$$

where  $\mu$  is the Milnor number at the origin. This number is well-defined because  $\mu(f_0) - \mu(f_s)$  is independent of  $s$  if  $s$  is sufficiently small, moreover by the upper semi-continuity of  $\mu$  this number is a non-negative integer.

The most famous result about the Milnor number and the topology of the family is Lê-Ramanujam's theorem [6]:

**Theorem 1.** *If  $n \neq 3$  and if  $\mu(f_0) = \mu(f_s)$  for all  $s \in [0, 1]$  then the topological types of  $f_0^{-1}(0)$  and  $f_s^{-1}(0)$  are equal.*

In other words, if the jump of the family  $(f_s)$  is 0 then  $f_0^{-1}(0)$  and  $f_s^{-1}(0)$  have the same topological type for sufficiently small  $s$ . Another motivation is that the jump of a family is crucial in the theory of singularities of polynomial maps at infinity.

The *jump*  $\lambda(f_0)$  of  $f_0$  is the minimum of the non-zero jumps of the  $(f_s)$  over all deformations of  $f_0$ . The problem, asked by N. A'Campo, is to compute  $\lambda(f_0)$ . We will only deal with plane curve singularities, that is to say  $n = 2$ .

As a corollary of our study we prove the following:

**Theorem 2.** *If  $f_0$  is an irreducible germ of plane curve and is Newton non-degenerate then*

$$\lambda(f_0) = 1.$$

A closely related question of V. Arnold [1] formulated with our definitions is to find all singularities with  $\lambda(f_0) = 1$ . S. Gusein-Zade [4] proved that

---

*Date:* February 1, 2008.

*2000 Mathematics Subject Classification.* 32G11, 14H20.

*Key words and phrases.* Deformation of singularity, Milnor number, Newton polygon.

there exist singularities with  $\lambda(f_0) > 1$  and as a corollary of a studied of the behaviour of the Milnor number in a deformation of a desingularization he proved Theorem 2 for all irreducible plane curves.

This note is organized as follows, in paragraphs 2 to 5 we define and calculate a weak form of the jump : the non-degenerate jump. In paragraph 6 we prove Theorem 2 and in paragraph 7 we give estimations when the germ is not irreducible. Finally in paragraph 8 we state some conjectures for the jump of  $x^p - y^q$ ,  $p, q \in \mathbb{N}$  and end by questions.

*Acknowledgments* : I am grateful to Norbert A'Campo for discussions.

## 2. KUŠNIRENKO'S FORMULA

We firstly recall some definitions (see [5]). Let  $f(x, y) = \sum_{(i,j) \in \mathbb{N}^2} a_{i,j} x^i y^j$  be an analytic germ of plane curve. Let  $\text{supp}(f) = \{(i, j) \in \mathbb{N}^2 \mid a_{i,j} \neq 0\}$  and  $\Gamma_+(f)$  be the convex closure of  $\bigcup_{(i,j) \in \text{supp}(f)} ((i, j) + \mathbb{R}_+^2)$  where  $(i, j) \in \text{supp}(f) \setminus \{(0, 0)\}$ . The *Newton polygon*  $\Gamma(f)$  is the union of the compact faces (called the *slopes*) of  $\Gamma_+(f)$ . We often identify a pair  $(i, j) \in \mathbb{N}^2$  with the monomial  $x^i y^j$ . Let  $f$  be *convenient* if  $\Gamma(f)$  intersects both  $x$ -axis and  $y$ -axis.

For a face  $\gamma$  of  $\Gamma(f)$ , let  $f_\gamma = \sum_{(i,j) \in \gamma} a_{i,j} x^i y^j$ . Then  $f$  is (*Newton*) *non-degenerate* if for all faces  $\gamma$  of  $\Gamma(f)$  the system

$$\frac{\partial f_\gamma}{\partial x}(x, y) = 0 ; \quad \frac{\partial f_\gamma}{\partial y}(x, y) = 0$$

has no solution in  $\mathbb{C}^* \times \mathbb{C}^*$ .

For a Newton polygon  $\Gamma(f)$ , let  $S$  be the area bounded by the polygon and  $a$  (resp.  $b$ ) the length of the intersection of  $\Gamma(f)$  with the axes  $x$ -axis (resp.  $y$ -axis). We set

$$\nu(f) = 2S - a - b + 1.$$

For a convenient germ  $f$  the local Milnor number verifies [5] :

### Theorem 3.

- $\mu(f) \geq \nu(f)$ ,
- if  $f$  is non-degenerate then  $\mu(f) = \nu(f)$ .

## 3. NON-DEGENERATE JUMP FOR CURVE SINGULARITIES

We will consider a weaker problem: Let  $f_0$  be a plane curve singularity and we suppose that  $(f_s)$  is a *non-degenerate deformation* that is to say for all  $s \in ]0, 1]$ ,  $f_s$  is Newton non-degenerate. The *non-degenerate jump*  $\lambda'(f_0)$  of  $f_0$  is the minimum of the non-zero jumps over all non-degenerate deformations of  $f_0$ . The new problem is to compute  $\lambda'(f_0)$ , in this note we explain how to compute it.

Obviously we have  $\lambda(f_0) \leq \lambda'(f_0)$  but this inequality can be strict. For example let  $f_0(x, y) = x^4 - y^4$ , then  $\lambda'(f_0) = 3$  which is obtained for the family  $f'_s(x, y) = x^4 - y^4 + sx^3$ . But  $\lambda(f_0) \leq 2$ , by the degenerate family  $f_s(x, y) = x^4 - (y^2 + sx)^2$  of jump 2.

## 4. COMPUTATION OF THE NON-DEGENERATE JUMP

For a convenient  $f_0$  there exists a finite set  $\mathcal{M}$  of monomials  $x^p y^q$  lying between the axes (in a large sense) and the Newton polygon  $\Gamma(f_0)$  (in a strict sense).

**Lemma 4.** *If  $f_0$  is non-degenerate and convenient then*

$$\lambda'(f_0) = \min_{x^p y^q \in \mathcal{M}} (\mu(f_0) - \mu(f_0 + s x^p y^q)),$$

for a sufficiently small  $s \neq 0$  (the minimum is over the non-zero values).

*Proof.* The proof is purely combinatoric and is inspired from [2]. For any polygon  $T$  of  $\mathbb{N} \times \mathbb{N}$ , we define as for  $\nu$  a number  $\tau(T) = 2S - a - b$ . Then  $\tau$  is additive : let  $T_1, T_2$  be polygons whose vertices are in  $\mathbb{N} \times \mathbb{N}$ , and such that  $T_1 \cap T_2$  has null area then  $\tau(T_1 \cup T_2) = \tau(T_1) + \tau(T_2)$ . By this additivity we can argue on triangles only. Moreover for a polygon  $T$  that do not contain  $(0, 0)$  we have  $\tau(T) \geq 0$ .

Now the jump for a non-degenerate family  $(f_s)$  corresponds to  $\tau(T)$  where  $T$  is the polygon “between”  $\Gamma(f_0)$  and  $\Gamma(f_s)$  ( $0 < s \ll 1$ ). Minimizing this jump is equivalent to minimizing  $\tau(T)$ . It is obtained for a polygon  $T$  for which all vertices except one are in  $\Gamma(f_0)$  and the last vertex is in  $\Gamma(f_s)$ . Then it is sufficient to add only one monomial corresponding to the latter vertex to obtain the required deformation.  $\square$

With this method we do not compute  $\mu(f_0)$ , nor  $\mu(f_0 + s x^p y^q)$  but directly the difference.

For a degenerate function  $f$  we denote by  $\tilde{f}$  a non-degenerate function such that  $f$  and  $\tilde{f}$  have the same Newton polygon:  $\Gamma(f_0) = \Gamma(\tilde{f}_0)$ . The non-degenerate jump for a degenerate function  $f_0$  can be computed with the easy next lemma:


**Lemma 5.** *Let  $f_0$  be degenerate.*

- $\lambda'(f_0) = \mu(f_0) - \mu(\tilde{f}_0)$  if  $\mu(f_0) - \mu(\tilde{f}_0) > 0$ ,
- else  $\lambda'(f_0) = \lambda'(\tilde{f}_0)$ .

## 5. AN EXAMPLE

For a given polynomial  $f_0$  it is very fast to see who will be the good candidates  $x^p y^q$  and hence to find  $\lambda'(f_0)$  after a very few calculus: we use that  $\mu(f_0) - \mu(f_0 + s x^p y^q) = \tau(T)$  where  $T$  is the zone between the Newton polygon of  $f_0$  and the one of  $f_0 + s x^p y^q$ .

For example let  $f_0(x, y) = x^4 - y^3$ . We draw its Newton polygon (see Figure 1). We easily see that the monomials  $x^p y^q$  that are candidates to minimize  $\tau$  for the zone between the Newton polygons are  $x^3$  (that will give a zone with  $\tau(T) = 2$ ) and  $xy^2$  that will give a zone with  $\tau(T) = 1$ . In that case the deformation will be  $f_s(x, y) = x^4 - y^3 + s x y^2$  and the jump of  $f_0$  is 1.

FIGURE 1. Example  $f_0(x, y) = x^4 - y^3$ 

## 6. IRREDUCIBLE CASE

In some cases we are able to give a formula for the computation of the jump. For example if  $f_0(x, y) = x^p - y^q$ , with  $\gcd(p, q) = d$  then by Bézout theorem there exists a pair  $(a, b)$  such that  $x^a y^b$  is in  $\mathcal{M}$  and such that the area  $T$  corresponding to the deformation  $f_s(x, y) = x^p - y^q + s x^a y^b$  is equal to  $d/2$ .

As an application we prove Theorem 2 cited in the introduction.

**Theorem 6.** *If  $f$  is irreducible and non-degenerate then  $\lambda(f) = \lambda'(f) = 1$ .*

*Proof.* We recall some facts from the book of Brieskorn-Knörrer [3, p.477]. For a germ of curve  $f$ , the number of slopes of a Newton polygon  $\Gamma(f)$  is lower or equal to the number  $r$  of irreducible components.

Moreover let  $R$  be the number of lattice points that belongs to  $\Gamma(f)$  minus 1. Then if  $f$  is non-degenerate we have  $R = r$ . The non-degenerate condition is not explicit in [3] but it is stated with an equivalent condition (a face is non-degenerate if and only if the corresponding polynomial  $g_i$  of [3, p.478] has only simple roots).

Then for an irreducible singular germ  $f$ ,  $\Gamma(f)$  has only one slope and  $f$  is convenient; moreover if  $f$  is non-degenerate then the extremities of  $\Gamma(f)$ , say  $x^p$  and  $y^q$ , verify  $\gcd(p, q) = 1$ . The non-degenerate jump of  $f$  is the same as for  $f_0 = x^p - y^q$  and is equal to 1 by Bézout theorem. Then  $\lambda'(f) = \lambda'(f_0) = 1$ , as  $0 < \lambda(f) \leq \lambda'(f) = 1$  it implies  $\lambda(f) = 1$ .  $\square$

## 7. NON IRREDUCIBLE CASE

More generally if  $f$  is convenient, non-degenerate, with one slope, let  $x^p, y^q$  be the extremities of the Newton polygon of  $f$ . Then  $f$  has the same non-degenerate jump as  $f_0 = x^p - y^q$ , we suppose  $p \geq q$  and we set  $d = \gcd(p, q)$ . The formula for  $\lambda'(f_0)$  is given by:

- (1) If  $1 \leq d < q \leq p$  then  $\lambda'(f_0) = d$  which is reached by a family  $f_s(x, y) = x^p - y^q + s x^a y^b$ ,  $a, b$  given by Bézout theorem.
- (2) If  $\gcd(p, q) = q$ , i.e.  $d = q$  then  $\lambda'(f_0) = q - 1$  which is reached with  $f_s(x, y) = x^p - y^q + s x^{p-1}$ .

We will give in paragraph 8 a conjectural value for  $\lambda(x^p - y^q)$ .

If there are several slopes with  $f$  convenient and non-degenerate then we can estimate  $\lambda'(f)$ . Let  $f = \prod_{i=1}^k f_i$  be the decomposition of  $f$  according to the slopes of  $\Gamma(f)$  (notice that  $f_i$  is not necessarily irreducible). If  $f_i$  is a smooth germ then we set (by convention)  $\lambda'(f_i) = 1$ . In fact  $f_i$  is smooth if and only if the corresponding slope  $\Gamma_i$  with extremities  $A_i, B_i$  verifies  $|x_{B_i} - x_{A_i}| = 1$  or  $|y_{B_i} - y_{A_i}| = 1$ . Then the following can be proved:

**Lemma 7.** *Let  $f$  be a convenient non-degenerate germ with several slopes, let  $f = \prod_{i=1}^k f_i$  be the decomposition according to the slopes.*

- (1) *If all the  $f_i$  are smooth then  $\lambda'(f) = 1$ .*
- (2) *If none of the  $f_i$  is smooth then*

$$\min_{i=1..k} \lambda'(f_i) \leq \lambda'(f) \leq \max_{i=1..k} \lambda'(f_i).$$

- (3) *In the other cases we have*

$$\min_{i=1..k} \lambda'(f_i) \leq \lambda'(f) \leq \max_{i=1..k} \lambda'(f_i) + 1.$$

We give some examples:

- (1) The family  $f_s(x, y) = (x + y^4)(x + y^2)(x^2 + y) + sy^4$  is of non-degenerate jump 1.
- (2) The family  $f_s(x, y) = (x^8 - y^6)(x^3 - y^2) + sxy^7$  gives  $\lambda'(f_0) = 2$  with  $\lambda'(x^8 - y^6) = 2$  and  $\lambda'(x^3 - y^2) = 1$ .
- (3) The family  $f_s(x, y) = (x^8 - y^6)(x^3 - y^2)(x^4 - y^4) + sx^5y^7$  verifies  $\lambda'(f_0) = 2$  while  $\lambda'(x^8 - y^6) = 2$  and  $\lambda'(x^3 - y^2) = 1$  and  $\lambda'(x^4 - y^4) = 3$ .
- (4) The family  $f_s(x, y) = (x + y^3)(x^4 + y^4)(x^2 + y) + sy^5$  verifies  $\lambda'(f_0) = 4$  with the smooth germs  $x + y^3, x^2 + y$  and  $\lambda'(x^4 + y^4) = 3$ .

## 8. CONJECTURES FOR THE JUMP

We give a conjectural value for  $\lambda(f_0)$  in the case that  $f_0 = x^p - y^q$  with  $p \geq q$ .

- (1) If  $\gcd(p, q) = 1$  then  $\lambda(f_0) = \lambda'(f_0) = 1$ .
- (2) If  $p = q$  and  $q$  is prime then  $\lambda'(f_0) = q - 1$ , with the family  $f_s(x, y) = x^q + y^q + sx^{q-1}$ . And we conjecture that  $\lambda(f_0) = q - 2$  with the family  $f_s(x, y) = x^q + y^q + s(x + y)^{q-1}$ .
- (3) If  $p = kq$  ( $k > 1$ ) and  $q$  is prime, then  $\lambda'(f_0) = q - 1$ , with the family  $f_s(x, y) = x^p + y^q + sx^{p-1}$ . It is conjectured that  $\lambda(f_0) = \lambda'(f_0)$ .
- (4) If  $q$  is not prime and  $p = kq, k \in \mathbb{N}^*$  then let  $q = ab$  with  $a \geq 2$  the smallest prime divisor of  $q$ . Then  $\lambda'(f_0) = q - 1 = ab - 1$  for the family  $f_s(x, y) = x^p - y^q + sx^{p-1}$ . It is conjectured that  $\lambda(f_0) = ab - b$ , which jump is reached for the family  $f_s(x, y) = x^p - (y^a + sx^{ka-1})^b$ .
- (5) If  $\gcd(p, q) = d$  with  $1 < d < q \leq p$  then  $\lambda'(f_0) = d$ . And it is conjectured that  $\lambda(f_0) = d$  too.

We make a remark for point (4), let  $g_0(x, y) = x^{p/b} - y^{q/b} = x^{ka} - y^a$ . Then  $g_0$  verifies the hypotheses of point (3) where we have conjectured  $\lambda(f_0) = a - 1$  for the deformation  $g_s(x, y) = x^{ka} - y^a - sx^{ka-1}$ . Then we calculate  $g_s(x, y)^b = (x^{ka} - y^a - sx^{ka-1})^b$  which is of course not a reduced polynomial. We develop and we have an approximation of  $g_s(x, y)^b$  if we set  $f_s(x, y) = x^{kab} - (y^a + sx^{ka-1})^b = x^p - (y^a + sx^{ka-1})^b$  with a jump equal to  $ab - b$ .

Apart from the conjectures above we ask some questions. Even if it seems hard to give a formula for the jump, maybe the following is easier:

*Question 1.* Find an algorithm that computes  $\lambda$ .

Finally the problem of the jump can be seen as a weak form of the problem of adjacency. For example the list of possible Milnor numbers arising from deformations of  $f_0(x, y) = x^4 - y^4$  is  $(9, 7, 6, 5, 4, 3, 2, 1, 0)$ . Then the gap between the first term  $9 = \mu(f_0)$  and the second term is the jump  $\lambda(f_0) = 2$ . Then the following question is a generalization of the problem of the jump.

*Question 2.* Give the list of all possible Milnor numbers arising from deformations of a germ.

#### REFERENCES

- [1] V. Arnold, Arnold's problems. *Springer-Verlag*, Berlin, 2004.
- [2] A. Bodin, Newton polygons and families of polynomials. *Manuscripta Math.* 113, (2004), 371–382.
- [3] E. Brieskorn and H. Knörrer, Plane algebraic curves. *Birkhäuser*, (1986).
- [4] S. Gusein-Zade, On singularities from which an  $A_1$  can be split off. *Funct. Anal. Appl.* 27 (1993), 57–59.
- [5] A. Kouchnirenko, Polyèdres de Newton et nombres de Milnor. *Invent. Math.* 32, (1976), 1–31.
- [6] Lê D. T. and C.P. Ramanujam, The invariance of Milnor's number implies the invariance of the topological type. *Amer. J. Math.* 98, (1976), 67–78.

LABORATOIRE PAUL PAINLEVÉ, MATHÉMATIQUES, UNIVERSITÉ DE LILLE I, 59655 VILLENEUVE D'ASCQ, FRANCE.

*E-mail address:* Arnaud.Bodin@math.univ-lille1.fr